

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/94855/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Awange, J., Forootan, Ehsan ORCID: <https://orcid.org/0000-0003-3055-041X>, Kusche, J., Kiema, J.B.K., Omondi, P., Heck, B., Fleming, K., Ohanya, S.O. and Goncalves, R.M. 2013. Understanding the decline of water storage across the Ramser-Lake Naivasha using satellite-based methods. *Advances in Water Resources* 60 , pp. 7-23. 10.1016/j.advwatres.2013.07.002 file

Publishers page: [http://www.sciencedirect.com/science/article/pii/S...](http://www.sciencedirect.com/science/article/pii/S0309170813001176)
<<http://www.sciencedirect.com/science/article/pii/S0309170813001176>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



This a short version of the paper

Understanding the decline of water storage across the Ramser-Lake Naivasha using satellite-based methods

Please cite: J. Awange; E. Forootan; J. Kusche; J.B.K. Kiema; P. Omondi; B. Heck; K. Fleming; S.O. Ohanya; R.M. Goncalves (2013). Understanding the decline of water storage across the Ramser-Lake Naivasha using satellite-based methods. *Advances in Water Resources*, Vol.60 , Pages 7–23, doi:10.1016/j.advwatres.2013.07.002

URL: <http://www.sciencedirect.com/science/article/pii/S0309170813001176>

Understanding the decline of water storage across the Ramser-Lake Naivasha using satellite-based methods

J.L. Awange^{a,e}, E. Forootan^b, J. Kusche^b, J.B.K. Kiema^c, P.A. Omondi^d, B.
Heck^e, K. Fleming^{a,f}, S.O. Ohanya^c, R. M. Gonçalves^g

^a*Western Australian Centre for Geodesy and The Institute for Geoscience Research, Curtin
University, Perth, Australia*

^b*Institute of Geodesy and Geoinformation, Bonn University, Bonn, Germany*

^c*Department of Geospatial and Space Technology, University of Nairobi, Nairobi, Kenya*

^d*IGAD Climate Prediction and Applications Centre (ICPAC), Nairobi, Kenya*

^e*Geodetic Institute, Karlsruhe Institute of Technology, Karlsruhe, Germany*

^f*Helmholtz Centre Potsdam, GFZ German Research Centre for Geosciences, Potsdam,
Germany*

^g*Department of Cartography Engineering, Federal University of Pernambuco (UFPE),
Geodetic Science and Technology of Geoinformation Post Graduation Program, Recife, PE,
Brazil*

Abstract

It has been postulated that Lake Naivasha, Kenya, has experienced a rapid decrease (and fluctuations) in its spatial extent and level between the years 2002 to 2010. Many factors have been advanced to explain this, with horticultural and floricultural activities, as well as climatic change, featuring prominently. This study offers a multi-disciplinary approach based on several different types of space-borne observations to look at the problem bedeviling Lake Naivasha, which is a Ramsar listed wetland of international importance. The data includes: (1) Gravity Recovery and Climate Experiment (GRACE) time-variable gravity field products to derive total water storage (TWS) variations within a region covering the Lakes Naivasha and Victoria basins; (2) precipitation records based on Tropical Rainfall Measurement Mission (TRMM) products to evaluate the impact of climate change; (3) satellite remote sensing (Landsat) images to map shoreline changes and to correlate these changes over time with possible causes; and (4) satellite altimetry observations to assess fluctuations in the lake's level. In addition, data from an in-situ tide gauge and rainfall stations as well as the output from the African Drought Monitor (ADM) model are used to evaluate the results. This study confirms that Lake Naivasha has been steadily declining with the situation being exacerbated from around the year 2000, with water levels falling at a rate of 10.2 cm/yr and a shrinkage in area of 1.04 km²/year. GRACE indicates that the catchment area of 4° × 4° that includes Lake Naivasha loses water at a rate of 1.6 cm/year for the period from August 2002 to May

Email addresses: J.Awange@curtin.edu.au (J.L. Awange), forootan@geod.uni-bonn.de (E. Forootan)

2006, and 1.4 cm/year for the longer period of May 2002 to 2010. Examining the ADM outputs also supports our results of GRACE. Between the time periods 2000-2006 and 2006-2010, the lake surface area decreased by 14.43% and 10.85%, respectively, with a corresponding drop in the water level of 192 cm and 138 cm, respectively, over the same periods. Our results show a correlation coefficient value of 0.68 between the quantity of flower production and the lake's level for the period 2002-2010 at 95% confidence level, indicating the probable impact of anthropogenic activities on the lake's level drop.

Key words: multi-disciplinary satellite data, lake hydrology, total water storage, Lake Naivasha, climate change, floriculture

1. Introduction

Lake Naivasha (Kenya, Figure 1) is the only freshwater lake in the Great Rift Valley of East Africa in an otherwise soda/saline lake series (Everard et al., 2002). In fact, it is the freshness of the water of Lake Naivasha that is the basis for its diverse ecology (Harper et al., 1990), and in 1995, it was declared as a Ramsar wetlands giving it an international status (see, e.g., Mekonnen et al. 2012). During the years 2002 to 2010, the lake has seen a rapid decline in its extent to the point where questions are being raised in the local media as to whether the lake is dying.

In the last decade, the level of Lake Naivasha has continued to drop with floriculture being blamed for excessive water extraction from the lake and aquifers, and the small holder farms in the upper catchment being blamed for nutrient loadings, leading to outcry in both the local and international media that this Ramsar site could be dying as a result of the very resource that it supports (see, e.g., ILEC, 2005; FWWCC, 2008; Mekonnen and Hoekstra, 2010). For example, Mekonnen and Hoekstra (2010) and Mekonnen et al. (2012) observed that the total virtual water exported in relation to the cut flower industry from the Lake Naivasha basin was 16 Mm³/yr during the period 1996-2005. This total virtual water (m³/yr) in relation to export cut flower and vegetables is obtained by multiplying the trade volumes (tones/yr) by their respective water foot print in Kenya (m³/ton), see e.g., Mekonnen et al., (2012). Other factors that have also been proposed as influencing Lake Naivasha's water changes include irregular rainfall patterns (Harper et al., 1990), and trade winds (Vincent et al., 1979). All of these discussions, therefore, point to the need for the reliable mapping of the lake and its basin in order to properly understand its dynamics.

Lack of reliable basin mapping techniques has hampered the proper monitoring of its changes, while also not allowing accurate predictions of the likely future situation, despite modelling methods being used to calculate its water balance (see e.g., Becht and Harper, 2002). The situation is compounded by the fact that Lake Naivasha has no surface outlet that could assist in hydrological monitoring, and that changes in its water level occur rapidly, over the order of several meters over just a few months, shifting the shoreline by several meters (Becht et al., 2005).

The emergence of satellite-based methods offers the possibility of providing a broader and more integrated analysis of the lake and its basin. Using time-variable gravity field products of the Gravity Recovery and Climate Experiment (GRACE) mission (Tapley et al., 2004), variations in the total water storage (TWS) of the region extending from the Lake Naivasha basin to Lake Victoria is assessed in this study, to determine whether the changes are climatic or human induced. GRACE-TWS products are then compared with soil moisture and separated into its compartments (i.e., precipitation and evaporation) using the African Drought Monitor (ADM) model. Changes in precipitation are further examined by analysing monthly products of the Tropical Rainfall Measurement Mission (TRMM), as well as four in-situ rainfall stations (Naivasha, Narok, Nakuru, and Kisumu), allowing us to determine the proportion of the fluctuations in Lake Naivasha that are related to changes in precipitation during a long-term period (1960 to 2010) and the study period (2002 to 2010). Note that analysing long-term precipitation variations also evaluates the impact of climate variability such as the dominant El Niño-Southern Oscillation (ENSO) phenomenon on the hydrological compartments of TWS variations within the region of study (Omondi et al., 2012; 2013a,b).

The fluctuations in the water level of Lake Naivasha are determined using both ground-based tide-gauge observations and satellite altimetry data (TOPEX/Poseidon and Jason-1). These results are then related to the use of satellite imagery (e.g., Landsat) and change detection techniques to map the shoreline changes of Lake Naivasha, analysing the trend of changes over the study period of interest, and correlating shoreline changes to the proposed causes. Therefore, this study pioneers the use of both space-borne and ground-based observations for monitoring Lake Naivasha.

2. Study Area

Lake Naivasha (00° 40' S - 00° 53' S, 36° 15' E - 36° 30' E) is the second largest fresh water lake in Kenya with a maximum depth of 8 m. It is situated in the Eastern African Rift Valley at an altitude of 1890 m above sea level and is approximately 80 km northwest of the Kenyan capital, Nairobi. Its basin (Figure 1) lies within the semi-arid belt of Kenya with mean annual rainfall varying from about 60 cm at the Naivasha township to some 170 cm along the slopes of the Nyandarua mountains, with open water evaporation estimated to be approximately 172 cm/year (Becht et al., 2005). Mount Kenya and the Nyandarua Range capture moisture from the monsoon winds, thereby casting a significant rain shadow over the Lake Naivasha basin (Becht et al., 2005). Unlike Lake Victoria which has its highest rainfall during the March-April-May (MAM) wet season (e.g., Awange et al., 2008a, b), the Lake Naivasha basin experiences its highest rainfall period during April-May-June (AMJ). There is also a short rainy season from October to November. The lake's levels, therefore, follow this seasonal pattern of rainfall cycle, with changes of several meters possible over a few months. Superimposed upon this seasonal behaviour are longer-term

77 trends, for example, there has been a change in the lake’s water level of 12 m
78 over the past 100 years (Becht et al., 2005).

FIGURE 1

79 The lake is fed by three main river systems: Gilgil, Malewa and Karati,
80 the last of which only flows during the wet season (see Figure 2). Becht et al.
81 (2005) observed that whereas a small portion of the groundwater evaporates and
82 escapes in the form of fumaroles in the geothermal areas, the remaining water
83 flows into Lakes Magadi and Elmentaita, taking thousands of years to reach
84 them. The basin’s water balance has been calculated from a model based upon
85 long-term meteorological observations of rainfall, evaporation and river inflows
86 (Becht and Harper, 2002). This model reproduced the observed level from 1932
87 to 1982 with an accuracy of 95% of the observed monthly level, differing by 0.52
88 m or less (ILEC, 2005). This pattern was, however, noticed to deviate after
89 1982 and by 1997, the differences had reached 3-4 m (Becht et al., 2005). In
90 fact, the onset of this reduced ability to model the lake’s level coincided with
91 the increase in horticultural and floricultural activities.

92 In general, three contemporary global water issues can be identified as occur-
93 ring in this region, namely *water scarcity/availability*, *water quality*, and *water*
94 *security*. While the focus of this study is on water scarcity/availability, several
95 previous works have focused on the problem of water quality and competition
96 for water resources within the study area (see e.g., Kitaka et. al 2002; Becht,
97 2007). Although water security issues are a reality in the Lake Naivasha basin,
98 few studies have been done to better understand the underlying conditions. For
99 example, Carolina (2002) asserts that the area of the Lake Naivasha basin is of
100 high economic and political importance to Kenya, and presents a wide variety
101 of economic activities based around the water resources, with many different
102 stakeholders often competing for the water resources.

103 The flower industry in Kenya has experienced a phenomenal growth, main-
104 taining an average growth rate of 20% per year over the last decade. It is an
105 industry that is the second largest export earner for Kenya, employing 50,000 -
106 60,000 people directly and 500,000 others indirectly through affiliated services
107 (KFC, 2011). Although flowers are now grown in many areas with temperate cli-
108 mate and an altitude above 1,500 m in Kenya, the region around Lake Naivasha
109 still remains the nation’s main floriculture farming center. The foremost cat-
110 egories of cut flowers exported from Kenya include roses, carnations, statice,
111 alstromeria, lilies and hypericum. Indeed, Kenya is arguably the largest ex-
112 porter for flowers in the world, supplying over 35% of cut flowers to the world’s
113 largest market - the European Union (KFC, 2011).

FIGURE 2

114 3. Datasets and Methodology

115 In Table 1, a complete set of data set used in this study are presented. Data
116 description is presented in our paper.

Table 1: Summary of the data sets used in this study.

Data	Period	Time steps
GRACE	2002.8 - 2010	monthly and 10-days
Altimetry	1992 - 2003	10-days
ADM	2002 - 2012	monthly
TRMM	2002 - 2012	monthly
In-situ rainfall	1960 - 2010	monthly
Tide gauge	1985 - 2010	monthly
Landsat	1989 - 2006	1989, 1995, 2000, and 2006
Flower export	1990 - 2010	yearly

FIGURE 3

4. Results and Discussions

4.1. Lake Level Analysis

The computed time series of level changes for Lake Naivasha derived from the T/P observations and in-situ measurements are shown in Figure 4. The T/P observation cover only the period between 1992 and 2003. The calculated satellite altimetry results were noisy at the first step, which may be related to the shallow depth of the lake (i.e., 8 m). To reduce this noise, the altimetry derived levels were smoothed using a moving average filter and interpolated according to the tide gauge time steps. As Figure 4 illustrates, the smoothed monthly altimetry derived levels are comparable to the available tide gauge measurements. We found a significant correlation coefficient of 0.69 between smoothed altimetry data and tide gauge observations (Figure 4,(Bottom)). Figure 4,(Top) confirms that although the lake level has been fluctuating both annually and seasonally over time up to around the year 2000, thereafter, a general downward trend at a rate of -10.2 cm/year before the onset of the 2007 ENSO rains is visible.

FIGURE 4

4.2. GRACE Analysis

Next, we estimated the changes in water mass over the Lake Naivasha basin as derived from GRACE observations. Because the GRACE-TWS results have a low spatial resolution, we compare two segments, one centred over Lake Victoria (to the west of Lake Naivasha) and the other centred over Lake Naivasha, as shown in Figure 5. The black boxes mark the areas where the GRACE-TWS and TRMM-total rainfall values were inferred. We chose a $4^\circ \times 4^\circ$ degree window as this is the limit to what can be confidently resolved from GRACE. Whereas GRACE is appropriate for areas the size of Lake Victoria (see section 3.1), our intention was to determine if it could still provide some information when comparing the variation of water within the basins of Lakes Naivasha and

143 Victoria, which in turn may be compared to TRMM data in order to infer the
 144 influence of climatic change to the region around Lake Naivasha.

FIGURE 5

145 Figure 6 (a) shows the TWS changes as described by the three GRACE
 146 products considered; CSR, CNES/GRGS and GFZ. Evidently, all GRACE so-
 147 lutions indicate water loss in both Lakes Naivasha and Victoria regions from
 148 2002 to late 2006. The increase in late 2006 is attributable to the ENSO effect,
 149 with water loss continuing again after an increase in late 2006-early 2007. Previ-
 150 ous studies have demonstrated that the fall in Lake Victoria during that period
 151 was due to anthropogenic factors such as the expanded Nalubale dam (see, e.g.,
 152 Awange et al., 2008a,b; Swenson and Wahr, 2009). Similar findings are shown
 153 by the cumulated water as illustrated in Figure 6 (b). The cumulative annual
 154 TWS of the Naivasha catchment lost water at a rate of 72 cm/yr from 2003 to
 155 May 2006. From January 2007 to December 2009, this loss was 41 cm/yr.

156 GRACE-TWS (as computed in Section 3) consists of a summation of ter-
 157 restrial water storage (WS), i.e., related to the catchment, and surface WS, i.e.,
 158 related to the lake itself. To enhance the interpretation of the GRACE's results
 159 in Figure 6, Lake Naivasha's surface WS changes are computed using its surface
 160 area, as shown with the solid-blue line in Figure 7 (top). To compute the blue
 161 line, the surface level changes (Figure 4) are transformed to the spherical har-
 162 monic domain and used to generate the surface WS changes time series (e.g.,
 163 as done in Swenson and Wahr (2009) for the derivation of hydrological trend of
 164 the East African lakes). The red line of Figure 7 (top) shows the time series
 165 computed for the GFZ GRACE products for the Naivasha region (i.e., Figure
 166 5; the right-hand-side box). Note that the leakage caused by Lake Victoria
 167 fluctuations is already removed from the red line, following Swenson and Wahr
 168 (2009). The catchment signal (terrestrial WS), shown on the bottom part of
 169 the figure as a black line, is the difference between GRACE-TWS (red line) and
 170 surface WS (blue line).

171 From Figure 7, we computed the slope of the blue line from August 2002
 172 to 2010 to determine the trend, obtaining a declining trend of 1.9 cm/year,
 173 while the period from May 2006 to 2010 saw a decline of 1.8 cm/year. After
 174 removing the signals of Lake Naivasha, the catchment area (black line in Figure
 175 7 (bottom)) loses water at a rate of 1.6 cm/year for the period from August 2002
 176 to May 2006 and 1.4 cm/year from May 2006 to 2010, thus signifying that not
 177 only is water lost from the Lake Naivasha but also from its catchment. The loss
 178 of water in the catchment could be attributed to floriculture and horticultural
 179 activities, and also boreholes providing water to the population that largely
 180 depends on the floricultural industry. In the next section, the use of ADM is
 181 employed to further enhance the GRACE results.

FIGURE 6

FIGURE 7

4.3. ADM Analysis

The red line in Figure 8 (top) shows the output of the ADM model derived from the right hand side of Eq. ?? compared to GFZ GRACE-TWS, averaged over the Lake Naivasha basin (i.e., Figure 7 (top), the red line). The mean of P for the years 2002 to 2012 was 103.7 mm and the standard deviation was 90.8 mm (maximum P was 645.8 mm). For E , ADM estimated a mean of 62.4 mm with a standard deviation of 20.1 mm (maximum E was 112.7 mm). The ADM-derived P and E are considerably smaller than what Becht et al. (2005) report, i.e., P of between 600 and 1700 mm/year and E of 1700 mm/year. Since the runoff parameter is not available after the year 2000 for Lake Naivasha (see also Ayenew and Becht, 2008) and the fact that Ojiambo et al. (2001) suggest that yearly R is negligible for the lake, we did not include it in our computations.

From the derived patterns, one can see that the ADM model responds more quickly to climatic variations such as ENSO in 2006 (red line in Figure 8 (A)) than the observed GRACE outputs (black line in Figure 8 (A)). Computing a correlation coefficient at 95% level of confidence shows a value of 0.68 between the two outputs, thus giving a reasonable level of agreement (Figure 8 (B)).

Visually comparing GRACE-derived terrestrial WS changes (shown by the black line in Figure 7 (Bottom)) with ADM-integrated soil moisture layers (Figure 8 (C)) reveals a similar pattern. The amplitude of the soil moisture signal is one third of the GRACE terrestrial WS changes. The reason for this inconsistency requires further research. Fitting a linear trend to the soil moisture results shows a TWS loss of 1.4 cm/year for the period from August 2002 to May 2006, and 0.6 cm/year from May 2006 to 2010.

Comparing the modeled precipitation (the green line in Figure 8 (D)) with in-situ precipitation (the cyan line in Figure 8 (D)), shows some inconsistencies, mainly in terms of the differences in the amplitudes between the modeled and in-situ values. A phase difference of one month is also evident between the two data sets. The dark-blue line in Figure 8 (D) represents the amount of evaporation changes for the period of July 2002 to 2012 showing almost steady range when compared to precipitations and soil moisture changes. As a result, one can see that the water capacity corresponding to soil moisture layers and rainfall is declining within the basin.

FIGURE 8

4.4. Rainfall Analysis

From the in-situ rainfall observations, the rainfall regime over the Naivasha basin has seen a downward trend since 1960 (see Figure 9). For instance, Figure 9 (bottom) shows a time series of the annual total (the black line), March-May (MAM, the blue line), June-August (JJA, the red line) and October-December (OND, the green line) rainfall seasons over Naivasha. In this study, we employed both graphical and statistical methods (described in WMO, 1966) to superficially test the significance of the observed trends (see also discussions in Wilks (1995) and Omondi et al. (2012; 2013a)). The data were analysed for

trends using linear regression, and the significance of trends was tested using the non-parametric Mann-Kendall tau test (Sneyers, 1990). An overview of the total amount of annual rainfall variation derived from the four stations is summarized in Figure 9, while their corresponding linear rates are reported on each graph. However, although the derived long-term linear trend values were negative, they were not large enough to pass the tau test (see also Omondi et al., 2013b).

There is also a high degree of variability, within both the wet periods (during strong El Niño years) and dry periods (during strong La Niña years). Several studies have investigated the relationship between eastern Africa rainfall and evolutionary phases of ENSO, and have shown strong relationship. Therefore, ENSO plays a significant role in determining the monthly and seasonal rainfall patterns in the East African region (e.g., Ogallo, 1988; Janowiak, 1988; Indeje, 2000; Mutemi, 2003, Nyakwada 2009 and Omondi et al., 2012). Considering the trends from the rain-gauge stations shown in Figure 9 suggests that the prolonged rainfall decrease over the catchment during the period 1960 to 2010 might contribute to the drop in the lake's level. Note that the linear trends for the period 2002-2010 (Figure 9, bottom) shows sharper decreasing values in all seasons and in the annual total rainfall than for 1960 to 2010. The result is in agreement with the variation in TWS as shown by GRACE analysis (Figure 7) and ADM (Figure 8). In Figure 11, the total amount of rainfall from the in-situ stations is compared to the GRACE TWS and the soil moisture WS from ADM.

FIGURE 9

Figure 10 illustrates the total rainfall of the catchment and its accumulated values as described by the TRMM 3B43 product over the 4×4 degree windows defined in Figure 5. The larger rainfall over Lake Victoria is seen both in terms of the time series, and also in the greater rate of increase in the accumulated values. Comparatively, while there seems to have been an increase in the precipitation rate over the Lake Victoria basin after late 2006, there seems to be little change in the rainfall over the Lake Naivasha basin. Comparing the TRMM results in Figure 10 with the GRACE results in Figure 6 for period 2002-2010, while no significant change is visible in the TRMM results, those from GRACE show a loss of water from the Lakes Naivasha and Victoria basins. This could therefore mean that the drop in Lake Naivasha's water level (as is the case for Lake Victoria) may be more influenced by anthropogenic factors compared to climatic factors.

FIGURE 10

4.5. Comparing TWS Changes Across Lake Naivasha with Rainfall

Figure 11 compares the accumulated annual TWS over the Naivasha catchment derived from GRACE as well as soil moisture from ADM (Figure 7) with the total annual rainfall variations derived from the four rainfall stations in Figure 9. To make the comparison easier, the values for 2003 are set to zero. As a

264 result, one can see that soil moisture and rainfall are decreasing between 2002
 265 to 2010. For 2006, GRACE still shows that TWS is decreasing, while precipita-
 266 tion increased as a result of ENSO, and soil moisture stays almost steady. The
 267 sharper rate of change that the GRACE results exhibit for 2002-2006 might
 268 also be related to the correlation of the derived TWS over Naivasha to that
 269 of Victoria. After 2006, again all component exhibit declining trends, showing
 270 that the impact of the 2006 ENSO has subsided.

FIGURE 11

271 4.6. Image Analysis

272 Next, we present the approach undertaken to map the shoreline variations
 273 of Lake Naivasha, using satellite images.

274 *Image classification:* To validate the results obtained from using GRACE
 275 and TRMM data sets, satellite remote sensing and GIS analysis were performed.
 276 Landsat imagery of the study area acquired from different epochs was employed
 277 and different land use / land cover types were discriminated. The interpreta-
 278 tion of Landsat imagery was undertaken using the minimum distance supervised
 279 classifier. The overall accuracy of the land use / land cover map was estimated
 280 to be 85.0% with a kappa statistic of 0.79. This meets the minimum thresh-
 281 old established by the United States Geological Survey (USGS) classification
 282 scheme (Anderson et al., 1976). As an example, the classification results for the
 283 first epoch (1989) are shown in Figure 12 (a). The classified image depicts a
 284 clear demarcation between land/vegetation and water, hence revealing a clear
 285 picture of the shoreline position. The red colour depicts water, yellow repre-
 286 sents general vegetation cover while green represent general bare land. There
 287 is a significant intrusion onto the northern shoreline by vegetation, indicating
 288 a positional change of the shoreline. Figure 12 (b), showing the second epoch
 289 examined (1995) shows a significant departure from Figure 12 (a), especially
 290 in the north, where vegetation has significantly receded, leaving only scattered
 291 traces in contrast to the 1989 image that showed a thick vegetation cover around
 292 the same area. There is also a change around Crescent Bay (formerly Crescent
 293 Island, see Figure 2). While the 1989 image shows a near excision of the bay
 294 from the main lake, the situation is different in the 1995 image. This is because
 295 of the general increase in water volume caused by increased rainfall over the
 296 same period.

297 Figure 12 (c) shows that there is an increase in water volume in 2000, due
 298 to more rainfall, compared with the preceding maps in Figures 12 (a and b).
 299 Crescent Bay has swollen with the south eastern section joined to the main lake
 300 to form the original Crescent Island, indicating an increase in water volume.
 301 This increase is probably due to the 1997 ENSO rainfall (see also the satellite
 302 altimetry results in Figure 4). The traces of vegetation that had infringed the
 303 northern part of the lake have fully disappeared by 2000. However, there are
 304 some traces of vegetation at the centre of the lake. These might be due to the
 305 presence of water lillies in the lake or traces of leaves transported by run-off

306 into the lake. Figure 12 (d) shows that the scattered traces of vegetation in
 307 the middle of the lake that were part of the preceding images have disappeared.
 308 However, the Crescent Bay has receded and a section of it is almost cut-off from
 309 the main lake to form an independent lake. There is also a significant change
 310 in the shape of the island when compared to the previous images. The amount
 311 of grassland cover has also increased along the shoreline compared to the 2000
 312 image, indicating a relationship between vegetation and the lake's surface area.

FIGURE 12

313 *Extraction:* It is visually clear from the classified land cover maps above
 314 that there is a perpetual shifting of the Lake Naivasha shoreline between dif-
 315 ferent epochs. Due to the difficulty in quantifying the amount and rate of the
 316 change, and in defining the actual trend through visual interpretation, the ac-
 317 tual position of the shoreline in each epoch was extracted and then compared
 318 to that obtained for the reference year, 1989. This allowed the actual change
 319 and subsequently the rate of change to be estimated.

320 This was done by digitizing the shoreline from the respective classified images
 321 in a Geographic Information System (GIS) environment using ArcGIS version
 322 9.3. The shorelines from each epoch were then overlaid to reveal the general
 323 change trend. To allow for a detailed analysis, the overlay result was further
 324 divided into five segments as shown in Figure 13. In general, the results show
 325 that there was an increase in water level in Lake Naivasha between 1989 and
 326 1995. This increase continued until 2000, however, the 2006 shoreline shows
 327 a decline in water level between 2000 and 2006. Detailed scrutiny shows that
 328 there was a steady northern (outward) shift of the shoreline from 1989 to the
 329 year 2000, indicating an increase in water level. This was followed by an inward
 330 shift in 2006, indicative of a drop in water level. However, the magnitude of the
 331 shoreline change is not uniform over the different epochs. The lack of uniformity
 332 can be attributed to variations in the local terrain, resulting in, obviously, the
 333 shoreline changing more in flatter terrain as opposed to steeper areas.

FIGURE 13

334 *Shoreline change and variation:* The surface area of the lake in each epoch
 335 was computed in the GIS environment. A summary of the changes in the surface
 336 area between different epochs for the different land cover classes, with 1989 as
 337 the reference year, is shown in Table 2, which indicates a direct relationship
 338 between the lake's level and surface area. The increase in area in 1995 compared
 339 to 1989 (i.e., 14.8%) is represented by an increase in water level (1.19 m). The
 340 situation is even more apparent between 1989 and 2000 where there is an increase
 341 of 20% in surface area and a corresponding increase of 2.33 m in water level.
 342 There was a drop in surface area of the lake between 2000 and 2006 (i.e., a drop
 343 of about 4.7%) and again this is shown by a drop of 1.92 m in the water level
 344 during this period. This general trend is corroborated by results obtained from
 345 both satellite altimetry and GRACE illustrated in Figures 4 and 7, respectively.

Table 2: Summary of the changes in the area of Lake Naivasha and the surrounding bare land and vegetation, with 1989 serving as the reference year (see Figures 12 and 14).

Year	Lake area (km ²)	Vegetation (km ²)	Bare land (km ²)	Mean Lake level (asl)	Lake area (%)	Vegetation (%)	Bare land (%)
1989	113.67	95.35	174.18	1885.41	0.00	0.00	0.00
1995	130.47	90.37	162.35	1886.60	14.78	-5.22	-6.89
2000	136.42	26.70	219.77	1887.74	20.01	-72.00	26.17
2006	130.07	48.44	204.12	1885.82	14.43	-49.20	17.19
2010	126.01	48.93	207.69	1884.44	10.85	-48.69	19.24

Figure 14 shows the variation of the surface area for the lake, vegetation, and bare land classes between 1989 and 2010.

FIGURE 14

From the above results, it is clear that Lake Naivasha has experienced shoreline variations over the last 17 years as indicated by the changes in surface area. There was a positive gain in area by 16.80 km² between 1989 and 1995 (i.e., 14.8%), with a further gain by the year 2000 of 5.95 km², due largely to the 1997 ENSO rainfall. However, there was a drastic decline in the surface area between 2000 and 2006, with the lake losing 6.35 km² of its surface area (i.e., 4.7%), indicating a recession in its shoreline. The surface area of the lake in 2006 is comparable to that of 1995 (both ~ 130 km²). After 2006, the lake continued shrinking with a surface area of 126.01 km² in 2010 (i.e., a reduction of about 7.6%). In general, from these images, it was calculated that the lake's area is shrinking at a rate of 1.04 km²/year. These findings agree with those of the satellite altimetry and tide gauge observations (see, e.g., Figure 4 in Sect. 4). The variation around the area shows that there is loss of vegetation around the lake as the lake surface area increases. There was a decline in the vegetation cover between 1989 and 1995, despite a gain in the surface area over this period. The same scenario was seen between 1995 and 2000. This can be attributed to the fact that the area around the lake is comprised of papyrus, which are normally swallowed by the increase in water level. There was, however, an increase in the vegetation cover between the years 2000 and 2006 as the lake receded and vegetation sprouted up along the shores of the lake .

4.7. Comparing Lake Levels with Rainfall and Flower Exports

In light of the previous results, the relationship between the decline of WS within the catchment, the lake itself, and the local and catchment precipitation were explored. To finalize this study, a simple comparison is made between the level of lake (one of the main water sources of the catchment) and other data sets considered in this work, including rainfall recorded by the Naivasha station (a representation of climate variability) and flower production (a representation of human use) (see Figure 15). Considering first rainfall and the lake's levels

(Figure 15 (A)), where we plot annual rainfall against annual average lake level, a correlation coefficient of -0.24 is obtained, suggesting no statistically significant correlation between these quantities. On the other hand, considering lake levels with flower production (Figure 15 (B)), where we have plotted tonnage of flower production against the annual averages of the lake levels for the years where the tonnage data were available, we find a strong statistically significant correlation, with a correlation coefficient of -0.68. This suggests strongly that flower exports could have influenced the reduction in Lake Naivasha's water level. Finally, Figure 15 (C) shows an insignificant correlation coefficient of -0.19 between rainfall and flower production.

Caution should be exercised, however, when one is interpreting the correlation results above. This is due to the fact that flower production, though it is a useful proxy for estimating water consumption in the Lake Naivasha region, and indeed constitutes the main cause of water consumption, depends on other factors unrelated to water withdrawal from the lake, e.g., in-put fertilizers. Therefore, an analysis of other factors that influence flower production, e.g., the amount of water withdrawn and used to irrigate the flowers, would be desirable. Along these lines, Mekonnen et al. (2012) quantified the water footprint within the Lake Naivasha Basin related to cut flowers and analysed the possibility of mitigating the footprint by involving cut-flower traders, retailers and overseas customers. Hagos (2008) assessed the possibility of using shallow and deep underground water, while Reta (2011) simulated a long term groundwater and lake water balance of Lake Naivasha in an attempt to establish the relationship between water consumption and water levels. Both Hagos (2008) and Reta (2011) highlighted the importance of underground water in the dynamics of Lake Naivasha's water levels. Such influence has been investigated, e.g., by Becht et al. (2002) and Becht and Nyaoro (2005), who considered the influence of groundwater fluctuations on Lake Naivasha and found it to have an important effect on the water balance of the Lake. In fact, Becht and Nyaoro (2005) deduced that the interaction of the groundwater and the Lake dynamics introduces a degree of inertia to the lake groundwater system, resulting in delayed reactions to external (meteorological) stresses where the groundwater acts as an extra reservoir absorbing water during wet periods and releasing water during droughts. Evaporation also plays a key role in Lake Naivasha's water balance as evident by the results of Farah et al., (2004), who obtained in-situ evaporation values at a grassland and woodland site in the Lake Naivasha basin for about a year. Another example of extraneous factors affecting Lake Naivasha's water levels is presented, e.g., in Olago et al. (2009), who showed that the hydrology of the Rift Valley is controlled mainly by climate and water table variation, among other factors.

FIGURE 15

5. Conclusions

As a Ramsar wetland, Lake Naivasha is a very important area not only to East Africa, but internationally. It supports a rich ecosystem with hundreds of species of diverse flora and fauna. Moreover, being the only freshwater lake in the Kenyan sector of the East African Rift, Lake Naivasha serves as the home of the flower industry in Kenya and is one of the most important flower producing regions world-wide. The results of this study have demonstrated that:

1. During the study period 1989 to 2010, Lake Naivasha experienced variation in its spatial extent and significant fluctuations in its level. However, from around the year 2000, a steady decline in its spatial extent has been observed with the lake receding at a rate of $1.41 \text{ km}^2/\text{year}$, accompanied by a corresponding drop in water level of about 33 cm/year.
2. Although the lake's level has been fluctuating both annually and seasonally over time in the past, there is a visible general downward trend observed from around 2000. This coincides with the period during which the flower exports from Kenya increased significantly. This is supported by the results of the linear regression analysis that gave a correlation coefficient of -0.68 between Lake Naivasha's water levels and the flower exports from the region for the period 2000-2010. Since much of the irrigation water used in the flower farms comes from Lake Naivasha, the recent decline in the lake water level and spatial extent could feasibly be largely attributed to adverse anthropogenic influences, with climatic factors such as prolonged rainfall decrease of the catchment during 1960-2010 also having a noticeable influence. A climatic influence is supported by the fact that in-situ rain gauge stations for the annual rainfall totals clearly indicate decreasing trends in the catchment area. The results support the findings in Mekonnen et al. (2012), who established a relationship between cut-flower production and level changes of Lake Naivasha.
3. Not only is the lake losing water, but also the catchment area of $4^\circ \times 4^\circ$ that includes Lake Naivasha as a whole is noticed to have lost water at a rate of 6.8 cm/yr from August 2002 to May 2008, and 1.7 cm/yr from May 2002 to 2010. The results are supported by the ADM output showing a decrease in soil moisture content, although the magnitude of the changes was one third of that shown by the GRACE results. While the long-term trend in the changes in precipitation was considerably less than those associated with soil moisture content and GRACE-TWS, the decline in the basin's water storage could possibly be related to the increased human use of groundwater within the catchment for horticulture, subsistence farming and domestic use.

These findings provide independent confirmation based on both ground and space-based observations on what has long been suspected, that is, floriculture has been exploiting the water resources of Lake Naivasha and the surrounding basin at an unsustainable rate. As pointed out in Sect. 4.7, however, floriculture may not be the sole cause of the decline of Lake Naivasha water levels.

Other factors, such as evaporation, fluctuation of groundwater level and climate among others, could also be contributing to the decline. Future studies on Lake Naivasha water levels should also include the effects of fluctuations of the Maleva and Gilgil rivers, especially the Maleva, which accounts for over 80% of inflows into the lake.

Remedial measures for the conservation and management of Lake Naivasha should thus be seriously considered before this Ramsar wetland becomes extinct. Already, the potential seriousness of the consequences arising from the decline of Lake Naivasha has finally been appreciated by the Government of Kenya, who has appointed an administrative body known as the Imarisha Lake Naivasha Management Board, for managing the Lake Naivasha Catchment Restoration Programme, whose aim is to restore Lake Naivasha and its catchment.

Acknowledgment

The authors are very grateful to the Editor Prof. G. Katul and the two anonymous reviewers for their comments, which helped to improve significantly the quality of the manuscript. J.L. Awange acknowledges the financial support of Alexander von Humboldt (Ludwig Leichhardt's Memorial Fellowship), The Institute for Geoscience Research (TIGeR), and Curtin Research Fellowship. He is grateful for the warm welcome and the conducive working atmosphere provided by his host Prof. Heck at the Geodetic Institute, Karlsruhe Institute of Technology (KIT). E. Foorotan and J. Kusche would like to thank the German Research Foundation (DFG), under the project BAYES-G, for the financial support. K. Fleming acknowledges the Australian Research Council's *Discovery Projects* funding scheme (project DP087738). The authors are grateful for the data sets used in this work, which were obtained from the following sources:

- Centre for Space Research (CSR) RL04 (<http://www.csr.utexas.edu/grace/asdp.html>).
- GFZ Information Systems and Data Center (<http://isdc.gfz-potsdam.de>).
- CNES/GRGS GRACE solutions (<http://bgi.cnes.fr:8110/geoid-variations/README.html>).
- TRMM, Goddard Earth Sciences Data and Information Services Center (<http://disc.sci.gsfc.nasa.gov/precipitation>).
- Landsat data, Global Land Cover Facility (GLCF), Maryland University, USA (<http://www.landcover.org/data/landsat/>).
- Tide-Gauge data. Lake Naivasha Riparian Association organization (<http://web.ncf.ca/es202/naivasha/>). Thanks to Mrs. Sara Higgins (the chair person of the Lake Naivasha Riparian Association) for providing these tide gauge data.
- Flower export data. Kenya Flower Council (<http://www.kenyafflowercouncil.org/>).

- 498 • Rainfall anomaly data. IGAD Climate Prediction and Application Centre,
499 Nairobi, Kenya.

500 References

- 501 Abiya, I. O., 1996. Towards sustainable utilization of Lake Naivasha, Kenya. *Lakes & Reservoirs: Research &*
502 *Management*, 2 (3-4), 231-242.
- 503 Adeyewa, Z.D., Nakamura, K., 2003. Validation of TRMM radar rainfall data over major climatic regions in major
504 climatic regions in Africa. *Journal of Applied Meteorology*, 42(2), 331-347.
- 505 Anderson, J.R., Hardy, E.E., Roach, J.T., Witmer W.E., 1976. A Land Use and Land Cover Classification System
506 for use with Remote Sensing Data, USGS Professional Paper 964, U.S. Geological Survey, Reston, Virginia,
507 4-38.
- 508 Awange, J.L., Ogalo, L., K-H. Bae, K-H., Were, P., Omondi, P., Omute, P., Omulo, M., 2008a. Falling Lake
509 Victoria levels: Is climate a contributing factor? *Climate Change*, 89: 281-297. doi:10.1007/s10584-008-
510 9409-x.
- 511 Awange, J., Sharifi, M., Ogonda, G., Wickert, J., Grafarend, E., Omulo, M., 2008b. The falling Lake Victoria water
512 level: GRACE, TRIMM and CHAMP satellite analysis of the lake basin. *Water Resource Management*, 22,
513 775-796. doi:http://dx.doi.org/10.1007/s11269-007-9191-y.
- 514 Awange, J., Ong'ang'a, O., 2006. Lake Victoria: ecology resource and environment. Springer, Berlin, 354 pp.
515 ISBN-13:978-3540325741.
- 516 Ayenew, T., Becht, R., 2008. Comparative assessment of the water balance and hydrology of selected Ethiopian
517 and Kenyan Rift Lakes. *Lakes & Reservoirs: Research & Management* 13(3), 181-196, doi:10.1111/j.1440-
518 1770.2008.00368.x.
- 519 Becht, R., 2007. Environmental effects of the floricultural industry on the Lake Naivasha basin. Project report,
520 International Institute for Geo-Information Science and Earth Observation (ITC), The Netherlands.
- 521 Becht, R., Harper, D.M., 2002. Towards an understanding of human impact upon the hydrology of Lake Naivasha,
522 Kenya. *Hydrobiologia*, 488: 1-12.
- 523 Becht, R., Odada, E.O., Higgins, S., 2005. Lake Naivasha: experience and lessons learned brief (Lake basin
524 management initiative): Experience and lessons learned briefs. including the final report: Managing lakes
525 and basins for sustainable use, a report for lake basin managers and stakeholders. Kusatsu: International
526 Lake Environment Committee Foundation (ILEC), 277-298.
- 527 Becht, R., Nyaoro, J. R., 2005. The influence of groundwater on lake-water management: the Naivasha case. In:
528 *Proceedings Volume II* (ed. by E. O. Odada, D. O. Olago, W. Ochola, M.Ntiba, S. Wandiga, K. Gichuki
529 & H. Oyieke) (Proc. 11th World Lakes Conference, Nairobi, Kenya, 31 October-4 November 2005), 384-388.
530 Ministry of Water and Irrigation, Kenya, and International Lake Environment Committee (ILEC), Japan.
531 http://www.ilec.or.jp/en/wp/wp-content/uploads/2013/02/World-Lake-Conference-11-Kenya-2005-Proceedings-
532 Volume-2.pdf
- 533 Becht, R., Odada, E.O., Higgins, S. 2005. Lake Naivasha : experience and lessons learned brief. In: Lake basin
534 management initiative : Experience and lessons learned briefs. including the final report: Managing lakes
535 and basins for sustainable use, a report for lake basin managers and stakeholders. Kusatsu : International
536 Lake Environment Committee Foundation (ILEC), 277-298.
- 537 Becker, M., Llovel, W., Cazenave, A., Grüntner, A. and Crétaux J.F., 2010. Recent hydrological behaviour of the
538 East African great lakes region inferred from GRACE, satellite altimetry and rainfall observations. *Comptes*
539 *Rendus Geoscience*, 342, 223-233. doi:10.1016/j.crte.2009.12.010.
- 540 Benada, J.R., 1997. TOPEX/POSEIDON User's Handbook. Technical report, Jet Propulsion Laboratory, Califor-
541 nia Institute of Technology Generation B (MGDR-B) Version 2.0.
- 542 Bettadpur, S., 2007. UTCSR Level-2 Processing Standards Document for Level-2 Product Release 0004. Center
543 for Space Research, The University of Texas at Austin, Rev 3.1, GRACE 327-742 (CSR-GR-03-03).
- 544 Bruinsma, S., Lemoine, J.M., Biancale, R., Valès, N., 2010. CNES/GRGS 10-day gravity field models (release 2)
545 and their evaluation. *Advances in Space Research*, 45(4), 587-601. doi:10.1016/j.asr.2009.10.012.
- 546 Carolina, B.F., 2002. Competition over water resources: analysis and mapping of water-related conflicts in the
547 catchment of Lake Naivasha (Kenya). MSc Thesis, International Institute for Geo-Information Science and
548 Earth Observation (ITC), The Netherlands.
- 549 Crétaux, J.F., Jelinski, W., Calmant, S., Kouraec, A., Vuglinski, V., Bergé-Nguyen, M., Gennero, M.-C., Nino,
550 F., Abarca Del Rio, R., Cazenave, A., Maisongrande, P., 2011. SOLS: A lake database to monitor in the
551 Near Real Time water level and storage variations from remote sensing data. *Advances in Space Research*,
552 47 1497-1507. doi:10.1016/j.asr.2011. 01.004.
- 553 Darling, W., Allen, D., Armannsson, H., 1990. Indirect detection of subsurface outflow from a rift valley lake.
554 *Journal of Hydrology*, 113 (1-4), 297-306. doi:http://dx.doi.org/10.1016/0022-1694(90)90180-6.
- 555 Everard, M., Harper, D.M., 2002. Towards the sustainability of the Lake Naivasha Ramsar site and its catchment.
556 *Hydrobiologia*, 488, (1-3), 191-203.
- 557 Everard, M., Vale, J.A., Harper, D.M., Tarras-Wahlberg. H., 2002. The physical attributes of the Lake Naivasha
558 catchment rivers. *Hydrobiologia*, 488, 13-25.
- 559 Farah H.O., Bastiaanssen, W.G.M., and Feddes, R.A., 2004. Evaluation of the temporal variability of the evapo-
560 rative fraction in a tropical watershed. *International Journal of Applied Earth Observation and Geoinfor-*
561 *mation*, 5, 129-140.

562 Flechtner, F., 2007. GFZ Level-2 Processing Standards Document for Level-2 Product Release 0004. Geo-
563 Forschungszentrum Potsdam, Rev. 1.0, GRACE 327-743 (GR-GFZ-STD-001).

564 Forootan, E., Awange, J., Kusche, J., Heck, B., Eicker, A., 2012. Independent patterns of water mass anomalies
565 over Australia from satellite data and models. *Remote Sensing of Environment*, 124, 427-443, dx.doi.org/10.1016/j.rse.2012.05.023.

566 FWWCC (Food & Water Watch and the Council of Canadians), 2008. Lake Naivasha withering under the as-
567 sault of international flower vendors. <http://www.canadians.org/water/documents/NaivashaReport08.pdf>
568 [Accessed on 25/08/2011].

569 Hagos G.G., 2008. Strategies to increase water use efficiencies of irrigated farm (south of Lake Naivasha, Kenya).
570 MSc. Thesis, Department of Geoinformation Science, ITC.

571 Harper, D.M., Mavuti, K.M., Muchiri, S.M., 1990. Ecology and management of Lake Naivasha, Kenya, in relation
572 to climatic change, alien species' introductions, and agricultural development. *Environmental Conservation*,
573 17 (04), 328-336.

574 Huffman, G.J., Adler, R.F., Bolvin, D.T., Gu, G., Nelkin, E.J., Bowman, K.P., Hong, Y., Stocker, E.F., Wolff,
575 D.B., 2007. The TRMM Multi-satellite Precipitation Analysis: Quasi-global, multi-year, combined-sensor
576 precipitation estimates at fine scale. *J. Hydrometeor.*, 8(1), 38- 55. doi:10.1175/JHM560.1.

577 ILEC (International Lake Environment Committee), 2005. Managing lakes and their basins for sustainable use.
578 A report for the lake basin managers and stakeholders. International Lakes Environmental Committee
579 Foundation: Kusatsu, Japan.

580 Indeje, M., Semazzi, F.H.M., Ogallo, L.J., 2000. ENSO signals in East African rainfall and their prediction
581 potentials. *Int. J. Climatol.* 20, 19-46.

582 Janowiak, J., 1988. An investigation of Interannual Rainfall Variability in Africa. *J. Clim.* 1, 240-255.

583 Jekeli, C., 1981. Alternative methods to smooth the Earth's gravity field. *Geodetic and Geoinformation Science*,
584 48. Department of Civil and Environmental Engineering and Geodetic Science, Ohio State University.

585 KFC (Kenya Flower Council), 2011. Kenya Flower Council. www.kenyaflowercouncil.org Accessed [25/7/2010].

586 Kitaka, N., Harper, D.M., Mavuti, K.M., 2002. Phosphorus inputs to Lake Naivasha, Kenya, from its catchment
587 and trophic state of the lake. *Hydrobiologia*, Vol. 488, 73-80. doi:10.1023/A:1023362027279.

588 Kummerow, C., Barnes, W., Kozu, T., Shiue, J., Simpson, J., 1998. The Tropical Rainfall Measuring Mission
589 (TRMM) sensor package. *Journal of Atmospheric and Oceanic Technology*, 15(3), 809-817.

590 Kummerow, C., Simpson, J., Thiele, O., Barnes, W., Chang, A., Stocker, E., Adler, R., Hou, A., Kakar, R., Wntz,
591 F., Aschroft, P., Kozu, T., Hing, Y., Okamoto, K., Iguchi, T., Kuroiwa, H., Im, E., Haddad, Z., Huffman,
592 G., Ferrier, B., Olson, W., Zipser, E., Smith, E., Wilhelm, T., North, G., Krishnamurti, T., Nakamura, K.,
593 2000. The status of the Tropical Rainfall Measuring Mission (TRMM) after two years in orbit. *Journal of*
594 *Applied Meteorology*, 39(12), 1965-1982.

595 Kusche, J., Schmidt, R., Petrovic, S., Rietbroek, R., 2009. Decorrelated GRACE time-variable gravity solutions
596 by GFZ, and their validation using a hydrological model, *Journal of Geodesy*, 83, 903-913. doi:10.1007/
597 /s00190-009-0308-3.

598 Luo, L., Sheffield, J., Wood, E.T., 2008. Towards a global drought monitoring and forecasting capability. *Sci-*
599 *ence and Technology Infusion Climate Bulletin*, 33rd NOAA Annual Climate Diagnostics and Prediction
600 Workshop Lincoln, NE, 20-24 October 2008.

601 Mather P., 2004. Computer processing of remotely sensed images. Third edition. 324 p. John Wiley & Sons, Ltd.
602 ISBN-13:978-0470849194.

603 Mekonnen, M.M., Hoekstra, A.Y., 2010. Mitigating the water footprint of export cut flowers from the Lake
604 Naivasha Basin, Kenya. *Value of Water Research Report Series No. 45*, UNESCO-IHE, Delft, The Nether-
605 lands.

606 Mekonnen, M.M., Hoekstra, A.Y., Becht, R., 2012. Mitigating the water footprint of export cut flowers from the
607 Lake Naivasha Basin, Kenya. *Water Resour Manage* 26, 3725-3742, doi: 10.1007/s11269-012-0099-9.

608 Mutemi, J.N., 2003. Climate anomalies over eastern Africa associated with various ENSO evolution phases. Ph.D.
609 Thesis, University of Nairobi, Kenya.

610 Nyakwada, W., 2009. Predictability of East African seasonal rainfall with sea surface temperature gradient modes.
611 Ph.D. Dissertation, Dept. of Met: University of Nairobi, Kenya.

612 Nicholson, S.E., 1998. Historical fluctuations of Lake Victoria and other lakes of the northern rift valley of
613 East Africa. In: *Environmental change and response in East African Lakes*, Kluwer Academic Publishers.
614 Lehman JT (Ed). 7-35. Kluwer Academic Publishers, Dordrecht, the Netherlands.

615 Nicholson, S., Some, B., McCollum, J., Nelkin, E., Klotter, D., Berte, Y., Diallo, B., Gaye, I., Kpabebe, G.,
616 Ndiaye, O., Noukpozoukou, J., Tanu, A., Thiam, A., Toure, A.A. and Traore, A., 2003. Validation of
617 TRMM and other rainfall estimates with a high-density gauge data set for West Africa: Part II: Validation
618 of TRMM rainfall products, *Journal of Applied Meteorology*, 42(10), 1355-1368.

619 Olago, D., Opere, A., and Barongo, J., 2009. Holocene palaeohydrology, groundwater and climate change in the
620 lake basins of the Central Kenya Rift, *Hydrological Sciences Journal*, 54:4, 765-780.

621 Ogallo, L.J., Janowiak, J.E., Halpert, M.S., 1988. Teleconnection between seasonal rainfall over Eastern Africa
622 and global sea surface temperature anomalies. *J. Meteor. Soc., Japan*, 66, 807-822.

623 Ojiambo, B.S., Poreda, R.J., Lyons, W.B., 2001. Ground Water/Surface Water Interactions in Lake Naivasha,
624 Kenya, Using $\delta^{18}\text{O}$, δD , and $^3\text{H}/^3\text{He}$ Age-Dating. *Groundwater* 39 (4), 526-533, doi:10.1111/j.1745-
625 6584.2001.tb02341.x.

626 Omondi, P., Awange, J.L., Ogallo, L.A., Okoola, R.A., Forootan, E., 2012. Decadal rainfall variability modes in
627 observed rainfall records over East Africa and their relations to historical sea surface temperature changes.
628 *J. Hydrol.* doi.org/10.1016/j.jhydrol.2012.07.003.

629 Omondi, P., Awange, J.L., Ogallo, L.A., Ininda, J., Forootan, E., 2013a. The influence of low frequency sea
630 surface temperature modes on delineated decadal rainfall zones in Eastern Africa region. *Advances in*
631 *Water Resources*, 54, 161-180. dx.doi.org/10.1016/j.advwatres.2013.01.001

632 Omondi, P., Awange, J.L., Forootan, E., et al. (2013b). Changes in temperature and precipitation extremes over
633 the Greater Horn of Africa region from 1961 to 2010. *International Journal of Climatology* (in press).

634 Reta G.L., 2011. Groundwater and lake water balance of lake Naivasha using 3-D transient groundwater model.
635 MSc. Thesis, Department of Geoinformation Science, ITC.

636 Richardson, J. L., Richardson E., 1972. History of an African rift lake and its climatic implications. *Ecological*
637 *Monographs*, 42(4): 499-534.

638 Rowlands, D., Luthcke, S., Klosko, F., Chinn, D., McCarthy, J., Cox, C., and Anderson, O., 2005. Resolving
639 mass flux at high spatial and temporal resolution using GRACE intersatellite measurements. *Geophysical*
640 *Research Letters*, 32(L04310). doi:10.1029/2004GL021908.

641 Sansome, H.W., 1952. The trend of rainfall in East Africa. *E. Afr. Meteorol. Dep., Tech. Mem.* 1. 14 p.

642 Sharifi, M., Forootan, E., Nikkhoo, M., Awange, J., Najafi, M. 2013. A point-wise least squares spectral analysis
643 (LSSA) of the Caspian Sea level fluctuations, using Topex/Poseidon and Jason-1 observations. *Journal of*
644 *Advances in Space Research*, 51 (1), 858-873. dx.doi.org/ 10.1016/j.asr.2012.10.001.

645 Sheffield, J., Goteti, G., Wood, E.F., 2006. Development of a 50-yr, high resolution global dataset of meteorological
646 forcings for land surface modeling. *J. Climate*, (13), 3088-3111. doi: http://dx.doi.org/10.1175/JCLI
647 3790.1.

648 Sheffield, J., Andreadis, K.M., Wood, E.F., Lettenmaier, D.P., 2009. Global and continental drought in the second
649 half of the 20th century: severity-area-duration analysis and temporal variability of large-scale events. *J.*
650 *Climate*, 22(8), 1962-1981. doi: http://dx.doi.org/10.1175/2008JCLI 2722.1.

651 Sneyers, R., 1990. On the statistical analysis of series of observations. WMO Technical Note, no. 143. ISBN-
652 13:978-9263104151.

653 Swenson, S., Wahr, J., Milly, P., 2003. Estimated accuracies of regional water storage variations inferred from
654 the Gravity Recovery and Climate Experiment (GRACE). *Water Resources Research*, 39(8). doi:10.1029/
655 2002WR001808.

656 Swenson, S., Wahr, J., 2006. Estimating large-scale precipitation minus evapotranspiration from GRACE satellite
657 gravity measurements. *J. Hydrometeorol*, 7 (2), 252-270, doi: http://dx.doi.org/10.1175/JHM478.1.

658 Swenson, S., Wahr, J., 2009. Monitoring the water balance of Lake Victoria, East Africa from space. *Journal of*
659 *Hydrology*, 370: 163-176. doi:http://dx.doi.org/10.1016/j.jhydrol.2009.03.008.

660 Tapley, B., Bettadpur, S., Watkins, M., Reigber, C., 2004. The gravity recovery and climate experiment: Mission
661 overview and early results. *Geophysical Research Letters*, 31. doi:10.1029/2004GL019920.

662 Tucker, C.J., Grant, D.M., Dykstra, J.D., 2004. NASA's global orthorectified Landsat data set. *Photogrametric*
663 *Engineering & Remote Sensing* 70(3), p. 313-322.

664 Vincent, C.E., Davies, T.D., Beresford, U.C., 1979. Recent changes in the level of Lake Naivasha, Kenya, as an
665 indicator of equatorial westerlies over East Africa. *Climatic Change* 2, 175-189.

666 Wahr, J., Molenaar, M., Bryan, F., 1998. Time variability of the Earth's gravity field: Hydrological and oceanic
667 effects and their possible detection using GRACE. *Journal of Geophysical Research* 103 (B12), 30205-30229.
668 doi:10.1029/98JB02844.

669 Wilks, D.S., 1995. Statistical methods in the atmospheric sciences. Academic press, pp 45-50, 399-402. ISBN-
670 13:978-0123850225.

671 WMO, 1966. Climate change. WMO Tech. Note no. 79, Geneva




Figure 1: Location map of the Lake Naivasha Basin (Becht et al., 2005).




Figure 2: Lake Naivasha drainage system (Becht et al., 2005).




Figure 3: Annual flower exports from Kenya (KFC 2011).




Figure 4: (Top) Time series of lake level height changes for Lake Naivasha as provided by satellite altimetry (T/P) and a tide gauge. (Bottom) Correlation between the lake level heights given by the tide gauge and the T/P altimetry.




Figure 5: The areas defined over the Lakes Naivasha and Victoria basins considered in the GRACE and TRMM analysis (see Figures 6 and 10). The red circle marks the location of Lake Naivasha.




Figure 6: Variations in stored waters (an integration of surface and terrestrial water storage changes) over Lakes Naivasha and Victoria derived from GRACE products. (a) Change in TWS and (b) accumulated changes of TWS in equivalent water volume (EWW) (see Figure 5, for the Victoria and Naivasha catchments).




Figure 7: Top, the red line shows the average TWS computed from the GFZ GRACE data (related to Figure 5, the black box on the right-side). The blue line is surface WS belonging only to Lake Naivasha. The catchment terrestrial WS signal is then obtained from the difference between GRACE-TWS signal (red line) and the Lake's surface WS signal (blue), i.e., the bottom graph with the black line.




Figure 8: (A) A comparison between the calculated TWS from the ADM TWS and the GRACE TWS, (B) shows the GRACE TWS against ADM TWS changes, (C) a basin averaged soil moisture layers over Naivasha, and (D) a comparison between model-derived precipitation and in-situ measurements along with the temporal pattern of evaporation.




Figure 9: Annual rainfall time series over four stations in the region of Lake Naivasha. From top, Nakuru (0.28°S, 36.1°E), Narok (1.1°S, 35.9°E), Kismu (0.1°S, 34.8°E) and Naivasha (0.72°S, 36.4°E) stations. For Naivasha, MAM [blue], JJA [red] and OND [green] yearly values are also provided.




Figure 10: Rainfall over the Lakes Naivasha and Victoria basins (see Figure 5) as provided by the TRMM 3B43 product. Rainfall amounts are shown by the solid lines and the accumulated values are dashed.




Figure 11: Comparing annual total water storage variations derived from GRACE with annual soil moisture contents (from ADM) and annual rainfall (from in-situ stations).




Figure 12: Surface-type classification results for the considered Landsat images. (a) 1989, (b) 1995, (c) 2000 and (d) 2006.




Figure 13: Segmentation of the changes in the Lake Naivasha shoreline for the years 1989, 1995, 2000 and 2006.




Figure 14: Variation in the area of the different land types around Lake Naivasha.




Figure 15: Comparing annual average lake levels with (A) rainfall observed at the Naivasha station and (B) flower exports. (C) Comparing annual average rainfall of the Naivasha station and flower exports. The solid lines are fitted linear trends, along with the correlation coefficients.